



**AWARENESS of PALLIATIVE CARE compared
among
HEALTHCARE PROFESSIONALS, PATIENTS and
CAREGIVERS
AN EXPERIENCE of FOUR DIFFERENT CENTRE**

Asst. Prof. Güliz DİRİMEN ARIKAN, MD. PhD.


Antalya, 20.09.2018

TOPICS of PRESENTATION


- Research Team
- Objectives
- Study Design
- Method
- Questionnaire
- Sampling
- Statistical Analysis




PROJECT TEAM



PROJECT TEAM



PROJECT TEAM



OBJECTIVES of THE PROJECT


PRIMARY OBJECTIVE

Demonstrating the awareness on palliative care of healthcare professionals, patients and their caregivers'

SECONDARY OBJECTIVES

- Demonstrating possible differences among professionals, patients and their caregivers' awareness on palliative care
- Planning the necessary interventions

STUDY DESIGN



METHOD

- ***The questionnaire design*** will be used to investigate the *demographical information and awareness on palliative care,*
- and *30 questions* will be asked.

Questions from the QUESTIONNAIRE:

Do you have religious belief?

Do you have a disease which is diagnosed and under the treatment?

Did you lose your relative because of a chronic disease?

Do you have a relative who is diagnosed with cancer?

Questions from the QUESTIONNAIRE:

Do you know about palliative care?

If you know about palliative care, where did you get the information?

-Hospital – Doctors – Nurses – TVs – Internet – ***My neighbour***

Questions from the QUESTIONNAIRE:

If you diagnosed with a fatal disease, do you want to be informed?

If you diagnosed with a fatal disease, whom you want to learn?

If you have a right to refuse CPR*, do you use it?

*CPR: Cardio Pulmonary Resuscitation

Some questions from the QUESTIONNAIRE:

- Do you know about palliative care?
- If you know about palliative care, where did you get the information?
 - Hospital – Doctors – Nurses – TVs – Internet – *My neighbour*
- If you diagnosed with a fatal disease, do you want to be informed?
- If you diagnosed with a fatal disease, whom you want to learn?
- If you have a right to refuse CPR*, do you use it?

***CPR**: Cardio Pulmonary Resuscitation

SAMPLING

Study Centres

- Anadolu Medical Centre Gebze Hospital,
- Dokuz Eylül University The Institute of Oncology, Medical Oncology Department
- Yeditepe University Teaching&Research Hospital,
- Yeditepe University Specialty Hospital

Participants

- * People who works at the centres,
- ***Healthcare professionals,***
- * People who seek care at the centres,
- ***Patients,***
- ***Caregivers.***

SAMPLING

- **P** (Patient): Patients, caregivers, healthcare professionals
- **I** (Intervention): No intervention – Questionnaire Design
- **C** (Comparison): Awareness comparison of three groups on palliative care
- **O** (Outcome): Awareness levels of groups
- **S** (Setting): Two cities, four centres – ISTANBUL, IZMIR
 - Anadolu Medical Centre Hospital,
 - Dokuz Eylül University The Institute of Oncology, Medical Oncology Department
 - Yeditepe University Teaching&Research Hospital,
 - Yeditepe University Specialty Hospital

SAMPLE SELECTION

- *Non-randomized purposive* sampling method will be used.



SAMPLE SIZE

Totally **1200** participants


* Sample size was calculated through,

- The number of healthcare professionals who included to the study and
- The number of patients and caregivers for each centre.

INCLUSION CRITERIA

- Turkish speaking and comprehending
- Age between 18-90
- Applied to one of the study centres as patient and/or caregiver
- Working as a healthcare professional at one of these centres:
 - Anadolu Medical Centre Hospital,
 - Dokuz Eylül University The Institute of Oncology, Medical Oncology Department
 - Yeditepe University Teaching&Research Hospital,
 - Yeditepe University Specialty Hospital

HEALTHCARE PROFESSIONALS WHO WERE PLANNED TO INCLUDE TO THE PROJECT



STATISTICAL ANALYSIS

- The information obtained from the questionnaire will be coded, entered, and analyzed using IBM SPSS version 22.
- Analysis will be done using descriptive statistics to obtain general characteristics of the study participants.
- Chi-square will be employed to determine the level of significance of groups of categorical variables such as gender, personnel cadre and duration of working.
- The p-values less than 0.05 will be considered as significant.





Thank you...

Asst. Prof. Gliz DİRİMEN ARIKAN, M.D. PhD.

guliz.dirimen@yeditepe.edu.tr

gulizdirimen@yahoo.co.uk